

Wavelet frames, Bergman spaces and Fourier transforms of Laguerre functions

Luís Daniel Abreu

ABSTRACT. The Fourier transforms of Laguerre functions play the same canonical role in wavelet analysis as do the Hermite functions in Gabor analysis. We will use them as analyzing wavelets in a similar way the Hermite functions were recently by Gröchenig and Lyubarskii in *Gabor frames with Hermite functions*, C. R. Acad. Sci. Paris, Ser. I 344 157-162 (2007). Building on Seip's work *Beurling type density theorems in the unit disc*, Invent. Math., **113**, 21-39 (1993), concerning sampling sequences on weighted Bergman spaces, we find a sufficient density condition for constructing frames by translations and dilations of the Fourier transform of the n th Laguerre function. As in Gröchenig-Lyubarskii theorem, the density increases with n , and in the special case of the hyperbolic lattice in the upper half plane it is given by

$$b \log a < \frac{4\pi}{2n + \alpha},$$

where α is the parameter of the Laguerre function.

1. Introduction

The "frame density problem" is one of the fundamental questions in applied harmonic analysis: how coarse we need to sample a continuous object so that the resulting discrete object is a frame in a certain Hilbert space? This question was sharply solved for frames in the Bargmann-Fock [8], [24], [25] and in the Bergmann space [28]. However, in the cases of general wavelet and Gabor transforms, very little is known and only a few very special windows and analysing wavelets are understood.

The short time Fourier (Gabor) transform with respect to a gaussian window can be written in terms of the Bargmann transform, mapping isometrically the space $L^2(\mathbf{R})$ onto the Bargmann-Fock space of entire functions. This is the reason why everything is known about the geometry of sequences that generate frames by sampling the Gabor transform with gaussian windows $g(t) = e^{-\pi t^2}$. Apart from this example, the only cases where a description is known of the lattice sequences that generate frames are the hyperbolic secant $g(t) = (\cosh at)^{-1}$ [18] and the characteristic function of an interval [19], which turned out to be a nontrivial problem. There is also a necessary density condition for Gabor frames due to Ramanathan and Steger [23] and it was recently observed that the frame property is stable under small perturbations of the lattice, given the window belongs to the Feichtinger algebra [10].

If we consider the wavelet case, things appear to be even more mysterious. Although the density and irregular grid problems have attracted some attention lately (see [1], [31], [22], [15] and references there in), there is no known counterpart of Ramanathan and Steeger theorem (there is, however, a version of the HAP property for wavelet systems [16]) and the only information available so far, on particular systems, concerns a special family of analyzing wavelets that maps the problem into Bergmann spaces: The wavelet transform, with positive dilation parameter, with respect to a wavelet of the form (Paul's Wavelet in some literature)

$$(1.1) \quad \psi_\alpha(t) = \left(\frac{1}{t+i} \right)^{\alpha+1},$$

can be rescaled as an isometrical integral transform between spaces of analytical functions, namely, between the Hardy space on the upper half plane $H^2(\mathbf{U})$ and the Bergman space in the upper half plane $A^{2\alpha+1}(\mathbf{U})$. It is natural to refer to this isomorphism as the Bergman transform (the designation analytic wavelet is also frequently used). Since the upper half plane can be mapped isomorphically onto the unit disc by using linear fractional transformations, we can construct a transform mapping the Hardy space onto the Bergman space in the unit disc.

The approaches used to deal with the special situations mentioned in the above paragraphs are based in techniques which differ from case to case. It seems highly desirable to follow a more structured approach. The natural place to look for this structure is within the context of Hilbert spaces of analytic functions, where the powerful methods from complex analysis may answer questions that seem hopeless otherwise. Following this line of reasoning implies using windows that allow to carry the problem to such spaces. In this direction a major step was taken recently by Göchenig and Lyubarskii [12], by considering Gabor systems with Hermite functions of order n as windows. They have proved that if the size of the lattice Λ is $< (n+1)^{-1}$ then the referred Gabor system is a frame and provided an example supporting their conjecture that the result is sharp.

In the Bargmann-Fock setting the Hermite functions play a very special role [11, pag. 57]. They are, up to normalization constants, pre-images, under the Bargmann transform, of the monomials $\{z^n\}$ and since the latter constitute an orthogonal basis of the Bargmann-Fock space, their pre-images also constitute an orthogonal basis of $L^2(\mathbf{R})$. That is, to the isomorphism

$$(1.2) \quad L^2(\mathbf{R}) \xrightarrow{\mathcal{B}} F^2(\mathbf{C})$$

corresponds

$$(1.3) \quad h_n \xrightarrow{\mathcal{B}} c_n z^n$$

where $\mathcal{B}$ is the Bargmann transform, $F^2(\mathbf{C})$ is the Bargmann-Fock space, h_n are the Hermite functions and c_n some constants dependent on n . They are canonical to time-frequency analysis in an additional sense, since they constitute the eigenfunctions of the time-frequency-localization operator with Gaussian window [3]. The Hermite functions are eigenfunctions of the Fourier transform and they can be used [20] to describe Feichtinger's algebra S_0 .

Wavelet and Gabor analysis share many similarities and many of their structural aspects can be bound together in a more general theory using representations of locally compact abelian groups [13], [9]. In looking for a Wavelet-analogue of Gröchenig-Lyubarskii structured approach to the density problem, we must first

clarify what functions should be used instead of the Hermite functions. In analogy to the above paragraph it is natural to consider the pre-images, under the Bergman transform, of the monomials $\{z^n\}$ in the unit disc (up to an isomorphism with the upper half plane). Since the $\{z^n\}$ form a basis of the weighted Bergman spaces in the unit disc (independently of the weight), their pre-images must be an orthogonal basis of the Hardy space $H^2(\mathbf{U})$ and it is reasonable to expect that such functions will play a similar role in Wavelet analysis as do the Hermite functions in Gabor analysis. Such functions are the Fourier transforms of the Laguerre functions and they will be ad-hoc denoted by S_n^α .

Additional structural evidence that the functions S_n^α are canonical in Wavelet analysis comes from the work of Daubechies and Paul [4], where it is shown that they are the eigenfunctions of a differential operator that commutes with a time-scale localization operator, once windows of the form (1.1) are chosen. This completely parallels the situation with the time-frequency-localization operators with gaussian windows, which commute with the harmonic oscillator and therefore have as eigenfunctions the Hermite functions [3]. It was observed by Seip [26] that these problems have a more natural formulation when mapped into the convenient spaces of analytical functions.

2. Description of the results

There exists an analogue of the correspondence between (1.2) and (1.3), but involving four different functional spaces and the corresponding bases: To the sequence of isomorphisms between the Hilbert spaces

$$(2.1) \quad L^2(\mathbf{0}, \infty) \xleftrightarrow{\mathcal{F}} H^2(\mathbf{U}) \xrightarrow{Ber^\alpha} A_{2\alpha+1}(\mathbf{U}) \xrightarrow{T_\alpha} A_{2\alpha+1}(\mathbf{D}),$$

where $A_{2\alpha+1}(\mathbf{U})$ and $A_{2\alpha+1}(\mathbf{D})$ are the weighted Bergmann spaces in the upper-half plane and in the unit disc, respectively, corresponds the relations between the basis of the respective spaces:

$$(2.2) \quad l_n^\alpha \xleftrightarrow{\mathcal{F}} S_n^\alpha \xrightarrow{Ber^\alpha} c_n^\alpha \Psi_n^\alpha \xrightarrow{T_\alpha} c_n^\alpha z^n.$$

for some constants c_n^α , where l_n^α is the n th Laguerre function of order n and parameter α , S_n^α is its Fourier transform and Ψ_n^α is a basis of $A_{2\alpha+1}(\mathbf{U})$ to be defined in section 4. This correspondence was implicit in the connection between papers [4] and [26] but since it was not stated explicitly, we devote section 4 to clarify how exactly does it work.

The functions S_n^α were computed recently in closed form [29], but the connection to the Wavelet transform seems to have been unnoticed. It is also shown in [29] that S_n^α are, up to a fractional transformation, defined in terms of a certain system of orthogonal polynomials on the unit circle. We will show that this family of polynomials is nothing more but the circular Jacobi orthogonal polynomials for which there is, for computational purposes, a very convenient three term recurrence formula. This connection will make our case to call the functions S_n^α the *rational Jacobi orthogonal functions*.

With a computational method available for evaluating the functions S_n^α and graphic evidence of their good localization properties (see for example the plots of their real versions in page 44 of [2]), it is natural to investigate how to use such functions as analysing wavelets and to obtain frames from the resulting discretization.

We will obtain the following sufficient condition on the density of the parameters of the hyperbolic lattice $\{(a^j b k, a^j)\}_{j,k \in \mathbf{Z}}$:

$$b \log a < \frac{4\pi}{2n + \alpha}$$

that is, as in Gröchenig and Lyubarskii result [12], this density increases with n . This will follow as a special case of the following more general theorem, which uses Seip's notion of lower Beurling density in the unit disk and constitutes the main result in this paper:

Theorem 5.1 *Let $\Gamma \subset D$ denote a separated sequence obtained from mapping the sequence $\{z_{m,j} = a_{m,j} + ib_{m,j}\} \subset U$ into the unit disk via a Cayley transform. If $D^-(\Gamma) > n + \frac{\alpha}{2}$ then $\{T_{a_{m,j}} D_{b_{m,j}} S_n^\alpha(\frac{t}{2})\}_{m,j}$ is a frame of $H^2(U)$.*

Case $\alpha = 0$ gives the sufficient part of theorem 7.1 in [28]. The proof of theorem 1 will be dramatically simplified with the observation of the following remarkable occurrence: There exist constants $\{C\}$ and sequences of numbers $\{a_k\}$, both depending on α and n , such that

$$(2.3) \quad S_n^\alpha\left(\frac{t}{2}\right) = C \sum_{k=0}^n a_k \psi_{k+\frac{\alpha}{2}}(t).$$

This will allow to obtain a formula which expresses the wavelet transform with window S_n^α as a linear combination of simple but non-analytic functions. As a result we will have to deal with a situation that is reminiscent of the non-analyticity problem that Gröchenig and Lyubarskii faced with formula (15) of [12] and which they solved by using the so called Wexler-Raz identities [11]. Since there is no known analogue of the Wexler-Raz identities in wavelet analysis, we have to deal with this in a more direct way. The key idea is to map the problem into the unit circle and use a deep sampling formula proved by Seip in [28]. Although this approach lacks the simplicity of [12] it gives the result for more general sequences, while the method in [12] works only for the lattice case due to the restrictions imposed by the use of the Wexler-Raz identities.

We organize our ideas as follows. The next section contains the main definitions and facts concerning wavelet transforms, Bergman spaces and Laguerre functions. Section four contains the evaluation of the pre-images required to build up the correspondences (2.1) and (2.2). The fifth section contains our main results on wavelet frames with Fourier transforms of Laguerre functions. We conclude the paper collecting some further properties of the functions S_n^α and clarifying their classification within known families of orthogonal polynomials.

3. Tools

3.1. The Bergman transform. Now we present a synthesis of ideas that appeared in the section 3.2 of [13] and in [4] (see also [2, pag. 31]) Here they will be exposed in such a way that the role of the Bergman spaces is emphasized.

Consider the dilation and translation operators

$$\begin{aligned} D_s f(x) &= |s|^{-\frac{1}{2}} f(s^{-1}x) \\ T_x f(t) &= f(t - x) \end{aligned}$$

and define

$$\psi^{x,s}(t) = T_x D_s \psi(t) = |s|^{-\frac{1}{2}} \psi\left(\frac{t-x}{s}\right).$$

The wavelet transform of a function f , with respect to the wavelet ψ is

$$W_\psi f(x, s) = \langle f, T_x D_s \psi(t) \rangle_{L^2(\mathbf{R})} = \int_{-\infty}^{\infty} f(t) \overline{\psi^{x,s}(t)} dt.$$

A function $\psi \in L^2(R)$ is said to be admissible if

$$\int_0^{\infty} |\mathcal{F}\psi(s)|^2 \frac{ds}{s} = K$$

where K is a constant. If ψ is admissible, then for all $f \in L^2(R)$ we have

$$(3.1) \quad \int_0^{\infty} \int_{-\infty}^{\infty} s^{-2} |W_\psi f(x, s)|^2 dx ds = K \|f\|^2$$

We will restrict ourselves to parameters $s > 0$ and functions $f \in H^2(\mathbf{U})$, where $H^2(\mathbf{U})$ is the Hardy space in the upper half plane $\mathbf{U} = \{z = x + is : s > 0\}$

$$H^2(\mathbf{U}) = \{f : f \text{ is analytic in } \mathbf{U} \text{ and } \sup_{0 < s < \infty} \int_{-\infty}^{\infty} |f(x + is)|^2 dx < \infty\}.$$

Let $\mathcal{F}$ denote the Fourier transform

$$(\mathcal{F}f)(t) = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} e^{-itx} f(x) dx$$

By the Paley-Wiener theorem, $H^2(\mathbf{U})$ is constituted by the functions whose Fourier transform is supported in $(0, +\infty)$ and belongs to $L^2(0, \infty)$.

Now take the special window $\psi_\alpha(t)$ defined in (1.1). Since

$$(3.2) \quad \overline{\mathcal{F}\psi_\alpha^{-x,s}(t)} = \mathbf{1}_{[0,\infty]} s^{\alpha+\frac{1}{2}} t^\alpha e^{i(x+is)t}$$

then

$$(3.3) \quad W_{\psi_\alpha} f(-x, s) = s^{\alpha+\frac{1}{2}} \int_0^{\infty} t^\alpha e^{izt} (\mathcal{F}f)(t) dt$$

where the function defined by the integral is analytic in $z = x + si$. The identity (3.1) gives

$$(3.4) \quad \int \int_{\mathbf{U}} |W_{\psi_\alpha} f(-x, s)|^2 s^{-2} dx ds = \|f\|_{H^2(\mathbf{R})}^2$$

This motivates the definition of the *Bergman transform*, or the *analytic wavelet transform*:

$$(3.5) \quad Ber^\alpha f(z) = \int_0^{\infty} t^\alpha e^{izt} (\mathcal{F}f)(t) dt = s^{-\alpha-\frac{1}{2}} W_{\psi_\alpha} f(-x, s),$$

where $z = x + is$ (see for instance [15], where the authors use this Bergman transform with the same normalization in the case $\alpha = 1$). Introducing the scale of weighted Bergman spaces

$$A_\alpha(\mathbf{U}) = \{f \text{ analytic in } \mathbf{U} \text{ such that } \int \int_{\mathbf{U}} |f(z)|^2 s^{\alpha-2} dx ds < \infty\},$$

it is clear from (3.4) and (3.5) that $Ber^\alpha f(z) \in A_{2\alpha+1}(\mathbf{U})$. We have therefore an isometric transformation

$$Ber^\alpha : H^2(\mathbf{R}) \rightarrow A_{2\alpha+1}(\mathbf{U})$$

The weighted Bergman spaces in the unit disc are denoted by $A_\alpha(\mathbf{D})$ and defined as

$$A_\alpha(\mathbf{D}) = \left\{ f \text{ analytic in } D \text{ such that } \int \int_D |f(z)|^2 (1 - |z|)^{\alpha-2} dx dy < \infty \right\}.$$

For detailed expositions of the theory of Bergman spaces we point out the monographs [32], [6], [14]

3.2. Frames and sampling sequences. A sequence of functions $\{e_j\}$ is said to be a frame in a Hilbert space H if there exist constants A and B such that

$$(3.6) \quad A \|f\|^2 \leq \sum_j |\langle f, e_j \rangle|^2 \leq B \|f\|^2.$$

In spaces of analytic functions a related concept to frames is the one of a sampling sequence. A set $\Gamma = \{z_j\}$ is said to be a sampling sequence for the Bergman space $A_\alpha(\mathbf{U})$ if there exist positive constants A and B such that

$$(3.7) \quad A \int \int_{\mathbf{U}} |f(z)|^2 y^{\alpha-2} dx ds \leq \sum_j |f(z_j)|^2 y_j^\alpha \leq B \int \int_{\mathbf{U}} |f(z)|^2 y^{\alpha-2} dx ds.$$

The corresponding definition of a sampling sequence for the Bergman space on the unit disc is

$$\begin{aligned} A \int \int_{\mathbf{D}} |f(z)|^2 (1 - |z|^2)^{\alpha-2} dz &\leq \sum_j |f(z_j)|^2 (1 - |z_j|^2)^\alpha \\ &\leq B \int \int_{\mathbf{D}} |f(z)|^2 (1 - |z|^2)^{\alpha-2} dz. \end{aligned}$$

Define the pseudohyperbolic metric on the unit disk by

$$\varrho(z, \zeta) = \left| \frac{z - \zeta}{1 - \bar{\zeta}z} \right|$$

Following [28], a sequence $\Gamma = \{z_j\} \subset \mathbf{D}$ is separated if $\inf_{n \neq j} \left| \frac{z_j - z_n}{z_j - \bar{z}_n} \right| > 0$ and its lower density D^- is given by

$$D^-(\Gamma) = \lim_{r \rightarrow 1} \inf_z \inf_z \frac{\sum_{\varrho(z_j, z) < r} (1 - \varrho(z_j, z))}{\log \frac{1}{1-r}}$$

The next Theorem is from [28]:

Theorem A *A separated sequence $\Gamma \subset \mathbf{D}$ is a sampling sequence for $A_\alpha(\mathbf{D})$ iff $D^-(\Gamma) > \frac{\alpha}{2} - \frac{1}{2}$.*

It is a generalization of an earlier lattice result [27]:

Theorem B *Let $\Gamma(a, b) = \{z_{jk}\}_{j,k \in \mathbf{Z}}$, where $z_{jk} = a^j(bk + i)$. $\Gamma(a, b)$ is a sampling sequence for $A_\alpha(\mathbf{U})$ if and only if $b \log a < \frac{4\pi}{\alpha-1}$.*

3.3. Fourier transforms of Laguerre functions. The Laguerre polynomials will play a central role in our discussion. One way to define them is by means of the Rodrigues formula

$$(3.8) \quad L_n^\alpha(x) = \frac{e^x x^{-\alpha}}{n!} \frac{d^n}{dx^n} [e^{-x} x^{\alpha+n}]$$

and this gives, in power series form

$$L_n^\alpha(x) = \frac{(\alpha+1)_n}{n!} \sum_{k=0}^n \frac{(-n)_k}{(\alpha+1)_k} \frac{x^k}{k!}.$$

For information on specific systems of orthogonal polynomials, we suggest [17]. The Laguerre functions are defined as

$$l_n^\alpha(x) = \mathbf{1}_{[0,\infty]}(x) e^{-x/2} x^{\alpha/2} L_n^\alpha(x)$$

and they are known to constitute an orthogonal basis for the space $L^2(0, \infty)$. By the Paley-Wiener theorem, the Fourier transform is an isomorphism between $H^2(\mathbf{U})$ and $L^2(0, \infty)$. Therefore, the Fourier transform of the functions l_n^α form an orthogonal basis for the space $H^2(\mathbf{U})$. From now on we will set

$$(3.9) \quad \mathcal{F}S_n^\alpha(t) = l_n^\alpha(t)$$

The functions $S_n^\alpha(t)$ can be evaluated explicitly using (3.8). This was already done by Shen in [29], where the author was interested in describing the orthogonal polynomials which arise from an application of the Fourier transform on the Laguerre polynomials. A description of this method of generating new families of orthogonal polynomials with an interesting historical account is in the paper [21] where a link is provided between Jacobi and Meixner-Polaczek polynomials. Following [29], we have

$$(3.10) \quad S_n^\alpha(t) = \frac{\Gamma\left(\frac{\alpha}{2} + 1\right) (1+\alpha)_n}{n!} \sum_{k=0}^n \frac{(-n)_k \left(\frac{\alpha}{2} + 1\right)_k}{k! (\alpha+1)_k} \left(\frac{1}{\frac{1}{2} - it}\right)^{k + \frac{\alpha}{2} + 1}$$

4. Bergman transform of $S_n^\alpha(t)$

In this section the correspondences (2.1) and (2.2) will be established via a direct calculation. The complex linear fractional transformations will play an important role, in a style that is reminiscent of the way they are used in the discrete series representation of $SL(2, \mathbf{R})$ over the Bergman space [7, chapter IX].

We first define a set of functions that, as we shall see later, constitute a basis of $A_\alpha(\mathbf{U})$. For every $n \geq 0$ and $\alpha > -1$ let

$$\Psi_n^\alpha(z) = \left(\frac{iz + \frac{1}{2}}{iz - \frac{1}{2}}\right)^n \left(iz - \frac{1}{2}\right)^{-\alpha-1}.$$

Now define a map T_α such that for every function $f \in A_\alpha(U)$ the action of T_α is

$$(4.1) \quad T_\alpha f(w) = f\left[\frac{i}{2} \frac{w+1}{w-1}\right] \left(\frac{1}{1-w}\right)^{\alpha+1}$$

The range space of T_α is a weighted Bergman space in the unit disc.

Lemma 4.1 *The map*

$$T_\alpha : A_\alpha(\mathbf{U}) \rightarrow A_\alpha(\mathbf{D})$$

is an unitary isometry between Hilbert spaces.

Proof. Argue as in the proof of lemma 1 in [7, pag.185]. $\square$

Proposition 4.1 *For $n = 0, 1, \dots$, the following relations hold:*

$$(4.2) \quad \text{Ber}^\alpha(S_n^{2\alpha}) = c_n^\alpha \Psi_n^\alpha, \text{ with } c_n^\alpha = (-1)^{\alpha+1} (2\alpha + n)!/n!$$

and, for $|z| < 1$,

$$(4.3) \quad T_\alpha(\Psi_n^\alpha) = z^n$$

In other words, the function $S_n^{2\alpha}$ is the pre-image, under $T_\alpha \circ \left(\frac{1}{c_n^\alpha} \text{Ber}^\alpha\right)$, of $z^n \in A_\alpha(D)$.

Proof. Using the definition of the Bergman transform (3.5) and (3.9) we have

$$\begin{aligned} \text{Ber}^\alpha S_n^{2\alpha}(z) &= \int_0^\infty t^\alpha e^{izt} (\mathcal{F} S_n^{2\alpha}(t))(t) dt \\ &= \int_0^\infty e^{(iz - \frac{1}{2})t} t^{2\alpha} L_n^{2\alpha}(t) dt \\ &= \frac{1}{n!} \mathcal{L} \left[\frac{d^n}{dx^n} [e^{-x} x^{2\alpha+n}] \right] \left(-iz + \frac{1}{2}\right) \\ &= c_n^\alpha \Psi_n^{2\alpha}(z), \end{aligned}$$

where in the last two identities we are applying Rodrigues formula (3.8) for the Laguerre polynomials and writing the integral in terms of the Laplace transform $\mathcal{L}$, whose well known properties establish the last identity. Now, the linear fractional transformation

$$w = \frac{iz + \frac{1}{2}}{iz - \frac{1}{2}}$$

is an analytic isomorphism between the upper half plane and the unit circle. Since the inverse of this transformation is given by

$$z = \frac{i}{2} \frac{w+1}{w-1},$$

a short calculation with the definition of T_α gives (4.3). $\square$

REMARK 1. $\{\Psi_n^\alpha(z)\}$ is a basis of $A_\alpha(U)$ and the map T_α is an unitary isomorphism between $A_\alpha(U)$ and $A_\alpha(\mathbf{D})$. Indeed, since $\{z^n\}$ is an orthogonal basis of the space $A_\alpha(\mathbf{D})$ [7, pag. 186] and it is contained in the range of T_α , T_α is onto and therefore an unitary isomorphism. The functions $\{\Psi_n^\alpha(z)\}$ form a basis for the space $A_\alpha(\mathbf{U})$ since they are the pre-images of the basis $\{z^n\}$.

As a consequence we obtain a new proof of the (known) isomorphic property of the Bergman transform.

COROLLARY 1. The transform $\text{Ber}^\alpha : H^2(\mathbf{R}) \rightarrow A_{2\alpha+1}(\mathbf{U})$ is an isometric isomorphism.

Proof. The isometry is a consequence of the isometric property of the wavelet transform, so we need only to prove that the $\text{Ber}^{\frac{\alpha}{2}}$ is onto. But in view of the preceding remark and Theorem 5.1, the range of $\text{Ber}^{\frac{\alpha}{2}}$ contains a basis of $A_\alpha(\mathbf{U})$. Therefore, $\text{Ber}^{\frac{\alpha}{2}}$ is onto. $\square$

REMARK 2. Similar calculations as we have seen here also play a role in [5], in the context of Laplace transformations and group representations and in [4], to obtain an explicit formula for the eigenvalues of the time-scale localization operator.

5. Wavelet frames with Fourier transforms of Laguerre functions

5.1. Preparation. We wish to construct wavelet frames with analysing wavelets S_n^α , by using the common discretization of the continuous wavelet transform via the hyperbolic lattice. Discretizing the scale parameter s in the Wavelet transform by a sequence a^j and the parameter x by $a^j b m$ gives

$$(5.1) \quad W_\psi f(a^j b m, a^j) = \langle f, T_{a^j b m} D_{a^j} \psi \rangle.$$

We want to know conditions in a and b under which $\{T_{a^j b m} D_{a^j} \psi\}$ is a frame, for a given wavelet ψ . More generally we can consider the non lattice problem replacing $(a^j b m, a^j)$ by more sequences (a_{jm}, b_{jm}) .

REMARK 3. Observe that it follows immediately from Theorem A and Theorem B that if $\Gamma \subset \mathbf{D}$ is the sequence obtained from mapping the sequence $\{z_{k,j} = a^j(bk + i)\} \subset \mathbf{U}$ into the unit disk via a Cayley transform, then $D^-(\Gamma) = \frac{2\pi}{b \log a}$ (to move between the notation of [28] and [27] set $\alpha = 2n + 1$).

REMARK 4. Observe that $\{T_{a^j b m} D_{a^j} \psi_\alpha\}_{j,k \in \mathbf{Z}}$ is a frame of $H^2(\mathbf{U})$ iff $\Gamma(a, b) = \{z = a^j b m + a^j i\}_{j,k \in \mathbf{Z}}$ is a set of sampling for $A_{2\alpha+1}(\mathbf{U})$. Indeed, since functions in $A_{2\alpha+1}(\mathbf{U})$ can be identified with Ber^α transforms of $H^2(\mathbf{U})$ functions, it follows from (3.7) and (??) that $\Gamma(a, b)$ is a set of sampling for $A_{2\alpha+1}(\mathbf{U})$ if and only if

$$A \int \int_{\mathbf{U}} |W_{\psi_\alpha} f(x, s)|^2 \frac{dx ds}{s^2} \leq \sum_{j,m} |W_{\psi_\alpha} f(a^j b m, a^j)|^2 \leq B \int \int_{\mathbf{U}} |W_{\psi_\alpha} f(x, s)|^2 \frac{dx ds}{s^2}.$$

Using (3.4) this is equivalent to

$$A \|f\|_{H^2(\mathbf{U})}^2 \leq \sum_{j,m} |\langle f, T_{a^j b m} D_{a^j} \psi_\alpha \rangle|^2 \leq B \|f\|_{H^2(\mathbf{U})}^2$$

which says that $\{T_{a^j b m} D_{a^j} \psi_\alpha\}_{j,k}$ is a frame in $H^2(\mathbf{U})$.

We now collect the required preliminary lemmas for the proof of the main result. The first was quoted in the introduction and is just a simple modification of the representation (3.10). The second appeared in [27] (see [6, pag. 160] for a proof) and the third one is a quite deep sampling theorem which is formula (30) in [28] with $s = 0$ and $\epsilon = 0$ (see also [6, pag. 216] for a direct derivation with this choice of parameters).

Lemma 5.2 The functions S_n^α can be written as the linear combination (2.3) of analyzing wavelets $\psi_{k+\frac{\alpha}{2}}(t)$ defined by

$$S_n^\alpha\left(\frac{t}{2}\right) = C \sum_{k=0}^n a_k \psi_{k+\frac{\alpha}{2}}(t),$$

where the constants C and the coefficients a_k given as

$$C = C^{\alpha,n} = \frac{\Gamma\left(\frac{\alpha}{2} + 1\right) (1 + \alpha)_n}{n!}$$

$$a_k = a_k^{\alpha,n} = (2i)^{k+\frac{\alpha}{2}+1} \frac{(-n)_k \left(\frac{\alpha}{2} + 1\right)_k}{k! (\alpha + 1)_k}.$$

Lemma 5.3 For $1 < s < t$ we have

$$(5.2) \quad \sum_j \frac{(1 - |z_j|^2)^s}{|1 - \bar{\zeta} z_j|^t} \leq C (1 - |\zeta|^2)^{s-t}$$

for any separated sequence $\{z_j\}$.

Lemma 5.4 *Let $D(\{w_j\}) > \beta$. Then every $f \in A_{2\beta+1}(D)$ satisfies*

$$(5.3) \quad (1 - |w|^2)^\beta f(w) = \sum_j (1 - |w_j|^2)^\beta f(w_j) h_j(w),$$

with the estimate for $h_j(w)$

$$(5.4) \quad |h_j(w)| \leq C \frac{(1 - |w|^2)(1 - |w_j|^2)}{|1 - \overline{w_j}w|^2}.$$

5.2. Main result. To prove our main result we will adhere to the following plan: we begin by deriving from Lemma 5.2 a formula expressing the wavelet transform with analysing wavelet S_n^α in terms of the wavelet transform with analysing wavelet $\psi_{k+\frac{\alpha}{2}}$. The proof of the right frame inequality is then straightforward (as usual in frame inequalities). To prove the left hand inequality we first rewrite the formula in terms of complex functions defined on the upper half plane and then map it into the unit disk by means of a Cayley transform. Manipulating the resulting formula and using lemma 5.3 and lemma 5.4 gives an estimate in the unit disk. We proceed backwards to the half plane to end up with the required estimate in terms of wavelet frames.

Theorem 5.1 *Let $\Gamma \subset D$ denote a separated sequence obtained from mapping the sequence $\{z_{m,j} = a_{m,j} + ib_{m,j}\} \subset U$ into the unit disk via a Cayley transform. If $D^-(\Gamma) > n + \frac{\alpha}{2}$ then $\{T_{a_{m,j}} D_{b_{m,j}} S_n^\alpha(\frac{t}{2})\}_{m,j}$ is a frame of $H^2(U)$.*

Proof. The definition of the wavelet transform gives, using (2.3),

$$\begin{aligned} W_{S_n^\alpha(\frac{t}{2})} f(-x, s) &= \left\langle f, T_{-x} D_s S_n^\alpha\left(\frac{t}{2}\right) \right\rangle \\ &= C \sum_{k=0}^n a_k \left\langle f, T_{-x} D_s \psi_{k+\frac{\alpha}{2}} \right\rangle \\ &= C \sum_{k=0}^n a_k W_{\psi_{k+\frac{\alpha}{2}}} f(-x, s). \end{aligned}$$

The right frame inequality is now easy: observe that, since $\{T_{a_{m,j}} D_{b_{m,j}} \psi_{k+\frac{\alpha}{2}}\}$ is a frame, then for every $k = 1, \dots, n$ there exists a B_k such that

$$\sum_{j,m} \left| \left\langle f, T_{a_{m,j}} D_{b_{m,j}} \psi_{k+\frac{\alpha}{2}} \right\rangle \right|^2 \leq B_k \|f\|_{H^2(U)}^2.$$

Let $B_n = \max_{1 \leq k \leq n} B_k$. Then,

$$\begin{aligned} \sum_{j,m} \left| \left\langle f, T_{a_{m,j}} D_{b_{m,j}} S_n^\alpha\left(\frac{t}{2}\right) \right\rangle \right|^2 &= \sum_{j,m} \left| C \sum_{k=0}^n a_k \left\langle f, T_{a_{m,j}} D_{b_{m,j}} \psi_{k+\frac{\alpha}{2}} \right\rangle \right|^2 \\ &\leq C \sum_{k=0}^n |a_k|^2 \sum_{j,m} \left| \left\langle f, T_{a_{m,j}} D_{b_{m,j}} \psi_{k+\frac{\alpha}{2}} \right\rangle \right|^2 \\ &\leq B \|f\|^2, \end{aligned}$$

where $B = B_n C \sum_{k=0}^n |a_k|^2$ and we are done.

Now the left frame inequality. Define the function $F(z)$ as

$$(5.5) \quad F(z) = W_{S_n^\alpha(\frac{1}{2})} f(-x, s) = C \sum_{k=0}^n a_k s^{k+\frac{\alpha}{2}+\frac{1}{2}} \text{Ber}^{k+\frac{\alpha}{2}} f(z)$$

where $z = x + is$ belongs to the upper half plane $\mathbf{U}$. We construct a related function in the unit disc, by setting $w = \frac{z-i}{z+i}$:

$$G(w) = C \sum_{k=0}^n a_k (1 - |w|^2)^{k+\frac{\alpha}{2}+\frac{1}{2}} G_{k+\frac{\alpha}{2}}(w)$$

where

$$G(w) = F\left(i \frac{1+w}{1-w}\right)$$

and

$$G_{k+\frac{\alpha}{2}}(w) = \left| \frac{2}{1-w} \right|^{2k+\alpha+1} \text{Ber}^{k+\frac{\alpha}{2}} f\left(i \frac{1+w}{1-w}\right) \in A_{2k+\alpha+1}(\mathbf{D}).$$

Therefore, if $D(\{w_j\}) > n + \frac{\alpha}{2}$ we can use (5.3) to write, for every $k \leq n$,

$$(1 - |w|^2)^{k+\frac{\alpha}{2}} G_{k+\frac{\alpha}{2}}(w) = \sum_j (1 - |w_j|^2)^{k+\frac{\alpha}{2}} G_{k+\frac{\alpha}{2}}(w_j) h_j(w).$$

As a result,

$$\begin{aligned} (1 - |w|^2)^{-\frac{1}{2}} G(w) &= C^{\alpha, n} \sum_{k=0}^n a_k (1 - |w|^2)^{k+\frac{\alpha}{2}} G_{k+\frac{\alpha}{2}}(w) \\ &= C \sum_{k=0}^n a_k \sum_j (1 - |w_j|^2)^{k+\frac{\alpha}{2}} G_{k+\frac{\alpha}{2}}(w_j) h_j(w) \\ &= \sum_j h_j(w) C \sum_{k=0}^n a_k (1 - |w_j|^2)^{k+\frac{\alpha}{2}} G_{k+\frac{\alpha}{2}}(w_j) \\ &= \sum_j h_j(w) (1 - |w_j|^2)^{-\frac{1}{2}} G(w_j). \end{aligned}$$

Now use estimate (5.4), then Cauchy-Schwarz and finally (5.2) in the following way.

$$\begin{aligned} (1 - |w|^2)^{-1} |G(w)|^2 &\leq \left[C \sum_j \frac{(1 - |w|^2)}{|1 - \overline{w_j} w|^2} (1 - |w_j|^2)^{\frac{1}{2}} G(w_j) \right]^2 \\ &\leq C \sum_j (1 - |w|^2)^2 |G(w_j)|^2 \sum_j \frac{(1 - |w_j|^2)^1}{|1 - \overline{w_j} w|^4} \\ &\leq C \sum_j (1 - |w|^2)^{-1} |G(w_j)|^2. \end{aligned}$$

Thus the factor $(1 - |w|^2)^{-1}$ cancels and we have

$$|G(w)|^2 \leq C \sum_j |G(w_j)|^2.$$

This last inequality gives at once

$$\int \int_D |G(w)|^2 (1 - |w|^2)^{-2} dw \leq C \sum_j |G(w_j)|^2$$

or, in the upper half plane,

$$A \int \int_{\mathbf{U}} |F(z)|^2 s^{-2} dx ds \leq \sum_j |F(z_j)|^2.$$

In the wavelet notation this is

$$A \int_0^\infty \int_{-\infty}^\infty \left| W_{S_n^\alpha(\frac{t}{2})} f(-x, s) \right|^2 s^{-2} dx ds \leq \sum_{j,m} \left| W_{S_n^\alpha(\frac{t}{2})} f(a^j b m, a^j) \right|^2.$$

Taking into account that $S_n^\alpha(\frac{t}{2})$ is admissible, we can apply (3.1) and (5.1) to write the above inequality as

$$(5.6) \quad A \|f\|^2 \leq \sum_{j,m} \left| \left\langle f, T_{a_m,j} D_{b_m,j} S_n^\alpha\left(\frac{t}{2}\right) \right\rangle \right|^2.$$

This is the left frame inequality. The sequence $\{T_{a_m,j} D_{b_m,j} S_n^\alpha(\frac{t}{2})\}_{j,m}$ is thus a frame. $\square$

In view of remark 5.1, the lattice result is a special case of our main theorem.

COROLLARY 2. *If $b \log a < \frac{4\pi}{2n+\alpha}$, then $\{T_{a^j b m} D_{a^j} S_n^\alpha(\frac{t}{2})\}_{j,m}$ is a frame of $H^2(\mathbf{U})$.*

REMARK 5. *Theorem 5.1 parallels theorem 3.1 in [12], which states that, in the lattice case of the time-frequency plane, if the density of the lattice Λ is $> n+1$ (or if the size of Λ is $< (n+1)^{-1}$) then the Gabor system $\{e^{2\pi i \lambda_2 t} H_n(t - \lambda_1) : \lambda = (\lambda_1, \lambda_2) \in \Lambda\}$, where H_n stands for the Hermite function of order n , is a frame for $L^2(\mathbf{R})$. In particular, if one is dealing with the Von Neumann lattice with parameters a and b , the condition is $ab < (n+1)^{-1}$. Corollary 5.2 makes this analogy even more explicit.*

REMARK 6. *It is interesting to notice that, since $Ber^\alpha f(z) = \int_0^\infty t^\alpha e^{izt} (\mathcal{F}f)(t) dt$, we clearly have*

$$i^k Ber^{k+\frac{\alpha}{2}} f(z) = \left(\frac{d}{dz}\right)^k Ber^{\frac{\alpha}{2}} f(z).$$

This allows to rewrite (5.5) as

$$W_{S_n^\alpha(\frac{t}{2})} f(-x, s) = C \sum_{k=0}^n \frac{a_k}{i^k} s^{k+\frac{\alpha}{2}+\frac{1}{2}} \left(\frac{d}{dz}\right)^k Ber^{\frac{\alpha}{2}} f(z)$$

which is, in some sense, reminiscent of Proposition 3.2 in [12].

REMARK 7. *Observe that combining the Paley-Wiener with the Plancherel theorem, we have $\|f\|_{H^2(\mathbf{U})} = \|f\|_{L^2(0,\infty)}$ and the results of Theorem 1 and 2 say also that we have a frame for $L^2(0,\infty)$.*

6. Further properties of the functions $S_n^\alpha(t)$

The notation

$$F(a, b; c; x) = \sum_{k=0}^{\infty} \frac{(a)_k (b)_k}{k! (c)_k} x^k.$$

for the hypergeometric function is used in this section. Rewriting S_n^α in this notation gives

$$S_n^\alpha(t) = C^{\alpha, n} \left(\frac{1}{2} - it \right)^{-\frac{\alpha}{2}-1} F\left(-n, \frac{\alpha}{2} + 1; \alpha + 1; \frac{1}{\frac{1}{2} - it}\right)$$

(observe that the infinite sum becomes a polynomial of order n , since $(-n)_k = 0$ if $k > 0$). Composing the functions $S_n^\alpha(t)$ with the fractional linear transformation

$$(6.1) \quad z = \frac{2t - i}{2t + i}$$

the result is

$$(6.2) \quad S_n^\alpha(t) = \Gamma\left(\frac{\alpha}{2} + 1\right) (1 - z)^{\frac{\alpha}{2}+1} g_n^\alpha(z)$$

where

$$g_n^\alpha(z) = \frac{(\alpha/2)_n}{n!} F\left(-n, \frac{\alpha}{2} + 1; -n - \alpha/2 + 1; z\right)$$

is a polynomial in z of degree n . This was pointed out in [29]. It was also shown that these polynomials satisfy the orthogonality

$$(6.3) \quad \int_{|z|=1} g_n^\alpha(z) \overline{g_m^\alpha(z)} |1 - z|^\alpha \frac{dz}{z} = 0 \text{ if } m \neq n$$

and therefore are orthogonal on the unit circle with respect to the weight

$$(6.4) \quad w(z) = \sin^\alpha \frac{\theta}{2} d\theta.$$

This fact implies many properties, since there exists a very rich theory for orthogonal polynomials on the unit circle (see [30] and references therein and also chapter 8 of [17]). For example, the general theory assures that all the zeros of $g_n^\alpha(z)$ lay within the unit disc.

REMARK 8. *Setting $a = \frac{\alpha}{w}$ in Example 8.2.5 at [17], and using the identity $1 - e^{i\theta} = 4 \sin^2 \frac{\theta}{2}$ to write the measure (8.2.21) as (6.4) we recognize that the polynomials $g_n^\alpha(z)$ are, up to a normalization, a family of orthogonal polynomials on the unit circle known as the circular Jacobi orthogonal polynomials.*

REMARK 9. *From (6.3) the functions $z^{-\frac{1}{2}}(1 - z)^{\frac{\alpha}{2}} g_n^\alpha(z)$ are orthogonal on the circle and form a basis of the Hardy space on the unit disc*

$$H^2(\mathbf{D}) = \{f : f \text{ is analytic in } \mathbf{D} \text{ and } \sup_{r < 1} \int_0^{2\pi} |f(re^{it})|^2 dt < \infty\}$$

it is also clear that $S_n^\alpha(t)$ are orthogonal on the real line (the boundary of the upper half plane). Since $g_n^\alpha(z)$ are the circular Jacobi orthogonal polynomials, the basis functions $z^{-\frac{1}{2}}(1 - z)^{\frac{\alpha}{2}} g_n^\alpha(z)$ are the circular Jacobi orthogonal functions and it is therefore natural to call $S_n^\alpha(t)$ the rational Jacobi orthogonal functions.

REMARK 10. From the general theory [17, (8.2.10)] follows that, if κ_n is the leading coefficient of the polynomial, then the sequence of polynomials $\{g_n^\alpha(z)\}$ satisfies a three term recurrence relation

$$\kappa_n g_n^\alpha(0) g_{n+1}^\alpha(z) + \kappa_{n-1} g_{n+1}^\alpha(0) z g_{n-1}^\alpha(z) = [\kappa_n g_{n+1}^\alpha(0) + \kappa_{n+1} g_n^\alpha(0) z] g_n^\alpha(z)$$

where κ_n is the leading coefficient of the polynomial. From the explicit representation of the polynomials $g_n^\alpha(z)$ it is easily seen that

$$\kappa_n = \frac{(\frac{\alpha}{2} + 1)_n}{n!}; \quad \phi_n(0) = \frac{\alpha(\frac{\alpha}{2} + 1)_{n-1}}{2n!}$$

This three term recurrence relation provides a effective method for computational purposes: To evaluate the functions $S_n^\alpha(t)$ it is sufficient to combine this recurrence relation with formulas (6.1) and (6.2).

References

- [1] A. Aldroubi, C. Cabrelli, U. M., Molter *Wavelets on irregular grids with arbitrary dilation matrices and frame atoms for $L^2(\mathbf{R}^d)$* , Appl. Comput. Harmon. Anal. 17, no. 2, 119–140 (2004).
- [2] I. Daubechies, "Ten lectures on wavelets", CBMS-NSF Regional conference series in applied mathematics (1992).
- [3] I. Daubechies, *Time-frequency localization operators: A geometric phase space approach* IEEE Trans. Inform. Theory, 34, pp. 605-612. (1988).
- [4] I. Daubechies, T. Paul, *Time-frequency localization operators: A geometric phase space approach II. The use of dilations and translations*. Inverse Prob., pp 661-680. (1988).
- [5] M. Davidson, G. Ólafsson, G. Zhang, *Laguerre polynomials, restriction principle, and holomorphic representations of $SL(2; \mathbf{R})$* . Acta Appl. Math. 71, no. 3, 261–277 (2002).
- [6] P. Duren, A. Schuster, "Bergman spaces", Mathematical Surveys and Monographs, 100. American Mathematical Society, Providence, RI, (2004).
- [7] S. Lang, " *$SL(2, \mathbf{R})$* ", Springer (1985).
- [8] Y. Lyubarskii, *Frames in the Bargmann space of entire functions*, Entire and subharmonic functions, 167-180, Adv. Soviet Math., 11, Amer. Math. Soc., Providence, RI (1992)
- [9] H. G. Feichtinger, K. Gröchenig, *Banach spaces related to integrable group representations and their atomic decompositions, I*, J. Funct. Anal. **86** (2), 307-340 (1989).
- [10] H. G. Feichtinger, N. Kaiblinger, *Varying the time-frequency lattice of Gabor frames*. Trans. Amer. Math. Soc. 356 (2004), no. 5, 2001–2023
- [11] K. Gröchenig, "Foundations of time-frequency analysis", Birkhäuser, Boston, (2001)
- [12] K. Gröchenig, Y. Lyubarskii, *Gabor frames with Hermite functions*, C. R. Acad. Sci. Paris, Ser. I 344 157-162 (2007).
- [13] A. Grossman, J. Morlet, T. Paul, *Transforms associated to square integrable group representations II: examples*, Ann. Inst. H. Poincaré Phys. Théor., **45**, 293-309, (1986).
- [14] H. Hedenmalm, B. Korenblum, K. Zhu, "Theory of Bergman spaces." Graduate Texts in Mathematics, 199. Springer-Verlag, New York, (2000).
- [15] C. Heil, G. Kutyniok, *Density of wavelet frames*, J. Geom. Anal. 13, no. 3, 480-493. (2003).
- [16] C. Heil, G. Kutyniok, *The Homogeneous Approximation Property for wavelet frames*, J. Approx. Theory., online first.
- [17] M. E. H. Ismail, "Classical and quantum orthogonal polynomials in one variable", Encyclopedia of Mathematics and its Applications No. 98. (2005).
- [18] A. J. E. M. Janssen; T. Strohmer, *Hyperbolic secants yield Gabor frames*. Appl. Comput. Harmon. Anal. 12, no. 2, 259–267 (2002).
- [19] A. J. E. M. Janssen, *Zak transforms with few zeros and the tie*, in "Advances in Gabor Analysis" (H.G. Feichtinger, T. Strohmer, eds.), Boston, 2003, pp. 31-70.
- [20] A. J. E. M. Janssen, *Hermite function description of Feichtinger's Space S_0* , J. Fourier Anal. Appl., **11** (5) 577-588 (2005).
- [21] H. T. Koelink, *On Jacobi and continuous Hahn polynomials*, Proc. Amer. Math. Soc. 124, no. 3, 887-898 (1996).

- [22] G. Kutyniok, *Affine density, frame bounds, and the admissibility condition for wavelet frames*. Constr. Approx., 25, no. 3, 239-253 , (2006)
- [23] J. Ramanathan, T. Steger, *Incompleteness of sparse coherent states*. Appl. Comput. Harmon. Anal. 2 , no. 2, 148–153 (1995).
- [24] K. Seip, *Density theorems for sampling and interpolation in the Bargmann-Fock space I*, J. Reine Angew. Math. 429, 91-106 (1992).
- [25] K. Seip, R. Wallstén, *Density theorems for sampling and interpolation in the Bargmann-Fock space II*, J. Reine Angew. Math. 429 (1992), 107-113.
- [26] K. Seip, *Reproducing formulas and double orthogonality in Bargmann and Bergman spaces*, SIAM J. Math. Anal. 22, No. 3 pp. 856-876 (1991).
- [27] K. Seip, *Regular sets of sampling and interpolation for weighted Bergman spaces*, Proc. Amer. Math. Soc. 117, no. 1, 213-220 (1993).
- [28] K. Seip, *Beurling type density theorems in the unit disc*, Invent. Math., **113**, 21-39 (1993).
- [29] L. C. Shen, *Orthogonal polynomials on the unit circle associated with the Laguerre polynomials*, Proc. Amer. Math. Soc. 129, no. 3, 873-879 (2001).
- [30] B. Simon, *OPUC on one foot*, Bull. Amer. Math. Soc. 42 (2005), 431-460.
- [31] W. Sun, X. Zhou, *Density and stability of wavelet frames*, Appl. Comput. Harmon. Anal. 15, no. 2, 117–133 (2003).
- [32] K. H. Zhu, *"Operator theory in function spaces"*. Monographs and Textbooks in Pure and Applied Mathematics, 139. Marcel Dekker, Inc., New York, 1990.